SCIENTIFIC REFERENCE GUIDE:

Air Pollutants and Health Impacts

+ lution

A comprehensive evidence-based guide to understanding what you breathe and how it affects your health.

Your health is in your air.

ABOUT THIS GUIDE

This scientific reference guide represents a comprehensive synthesis of environmental health research spanning five decades of epidemiological studies, controlled human exposure experiments, and mechanistic investigations.

The evidence base includes:

- **2,000+ peer-reviewed publications** from top-tier journals
- **50+ landmark cohort studies** tracking millions of participants
- **30+ systematic reviews and meta-analyses** pooling global evidence
- **EPA and WHO Integrated Science Assessments** representing scientific consensus

The health effects quantified here are not theoretical—they represent measurable, reproducible impacts demonstrated across diverse populations worldwide. Every percentage increase in mortality, every IQ point lost, and every disease risk elevation has been validated through multiple independent research teams using different methodologies in different populations, yet reaching remarkably consistent conclusions.

This convergence of evidence from molecular mechanisms to population-level outcomes establishes air quality as one of the most powerful, modifiable determinants of human health in the modern era.

This reference guide synthesizes peer-reviewed research from leading journals including NEJM, JAMA, The Lancet, Environmental Health Perspectives, and integrated science assessments from EPA and WHO. All statistics represent meta-analyses or large cohort studies with appropriate confidence intervals. For complete citations and methodological details, refer to the full scientific bibliography.

Last Updated: November 2025
Scientific Advisory: Lution Labs Clinical Research Team

Pristine air represents the evolutionary baseline —what human biology was designed to breathe. The gap between current exposure and pristine conditions represents your optimization opportunity:

- **PM2.5:** 8.2 μg/m³ gap (10.2 current vs. 2.0 pristine)
- **Ozone:** 22-27 ppb gap
- **NO_2:** 12-13 ppb gap
- **Lead:** 100-fold elevation in blood levels

UNITS EXPLAINED

Concentration Measures

- **μg/m³** = Mass per volume of air
- **ppb/ppm** = Parts per billion/million
- **CFU/m³** = Living organisms
- **EU/m³** = Endotoxin activity

Key Takeaways from the Science



1. **No Safe Thresholds**: PM2.5, lead, and benzene show health effects at the lowest measurable levels
2. **Indoor Often Worse**: NO_2, benzene, CO, and biological contaminants frequently exceed outdoor levels
3. **Cumulative Effects**: Multiple pollutants interact synergistically, amplifying health impacts
4. **Chronic Inflammation**: poor air quality keeps the immune system in a state of near-constant activation, with pernicious effects on virtually all biological systems
5. **Reversible Damage**: Studies prove reducing exposure rapidly improves health outcomes
6. **Vulnerable Populations**: Children, elderly, and those with preexisting conditions face amplified risks

OUTDOOR AIR POLLUTANTS:

Regulatory Standards and Health Science

POLLUTANT	TYPICAL LEVELS	MAJOR SOURCES
PM2.5 (Particulate Matter <2.5μm)	**Average USA:** 10.2 μg/m³ **EPA Standard:** 9.0 μg/m³ **WHO Guideline:** 5.0 μg/m³ **Pristine Air:** 2.0 μg/m³ *Indoor often 2-5× outdoor levels*	• Vehicle exhaust (especially diesel) • Power plant emissions • Industrial processes • Wildfires • **Indoor:** Cooking (gas stoves), candles, tobacco smoke
Ozone (O_3)	**Average USA:** 42 ppb **EPA Standard:** 70 ppb **WHO Guideline:** 51 ppb **Pristine Air:** 15-20 ppb *Summer peaks, formed photochemically*	• VOCs + NOx + sunlight • Vehicle emissions • Industrial facilities • Power plants • **Note:** Secondary pollutant - not directly emitted
Nitrogen Dioxide (NO_2)	**Average USA:** 15 ppb **EPA Standard:** 53 ppb **WHO Guideline:** 10 ppb **Pristine Air:** 2-3 ppb *Indoor can exceed 200 ppb during cooking*	• Motor vehicles (40-50% urban) • Power plants • **Indoor:** Gas stoves, unvented heaters • Industrial combustion
Sulfur Dioxide (SO_2)	**Average USA:** 8 ppb **EPA Standard:** 75 ppb **WHO Guideline:** 15 ppb **Pristine Air:** <1 ppb *Declined 70% since 2011*	• Coal-fired power plants • Industrial facilities • Metal processing • Volcanic activity • Marine vessel fuel
Carbon Monoxide (CO)	**Average USA:** 0.8 ppm **EPA Standard:** 9 ppm **WHO Guideline:** 3.5 ppm **Pristine Air:** 0.1-0.2 ppm *Indoor often exceeds outdoor*	• Vehicle exhaust • **Indoor:** Gas appliances, attached garages, tobacco smoke • Wood-burning stoves • Malfunctioning heaters
Benzene	**Average USA:** 0.5 μg/m³ **EPA:** No safe standard **WHO Guideline:** 1.7 μg/m³ **Pristine Air:** <0.1 μg/m³ *Indoor 2-10× outdoor*	• Traffic (84% of exposure) • **Indoor:** Cigarette smoke (primary), paints, adhesives • Gasoline vapors • Attached garages
Lead	**Average USA:** 0.01 μg/m³ **EPA Standard:** 0.15 μg/m³ **WHO Guideline:** 0.5 μg/m³ **Pristine Air:** <0.001 μg/m³ *Blood lead: 1-2 μg/dL typical vs. 0.016 pristine*	• Aviation fuel (largest current • Lead battery recycling • Industrial emissions • **Legacy:** Old paint, contaminated soil

HEALTH EFFECTS	LANDMARK SCIENTIFIC STUDIES
Mortality: 16% increase per 10 μg/m³ **Heart Attack:** +23% risk **Stroke:** +24% risk **Lung Cancer:** +9% risk **Dementia:** +14-40% risk **Life Expectancy:** 2-3 years lost in polluted cities **Cognitive:** Reduced brain volume, impaired memory	**Harvard Six Cities Study:** Established PM2.5-mortality link in 8,111 adults **American Cancer Society Study:** 138 adults confirmed cardiovascular mortality **Global Meta-Analysis** 107 cohorts showing effects below 10 μg/m³ **Dementia Study** 28 cohorts linking PM2.5 to cognitive decline
All-cause Mortality: +1.4% per 10 ppb **Respiratory Death:** +2.5% **COPD Death:** +5.6% **Heart Failure Death:** +7.4% **Lung Function:** Rapid decline **Asthma:** Attacks triggered at 70 ppb **Exercise Performance:** Reduced	**Cancer Prevention Study II:** 669,046 participants showing respiratory mortality **Meta-Analysis:** 99.9 million participants, established mortality risks **Controlled Exposure Studies:** Lung function decrements at current EPA standard
All-cause Mortality: +6% per 10 ppb **Cardiovascular Death:** +11% **Childhood Asthma:** +32% from gas cooking **Current Wheeze:** +15% **Respiratory Infections:** Increased susceptibility **Inflammation:** IL-6, IL-1β elevation	**Global Meta-Analysis:** 34 studies, 32 cohorts establishing mortality **Indoor NO_2 Study:** Gas cooking and childhood asthma link **Controlled Exposure:** Inflammatory response at 1.5 ppm
All-cause Mortality: +0.59% per 10 μg/m³ **Respiratory Death:** +0.67% **Asthma Attacks:** Triggered at 0.25 ppm **Airway Resistance:** +322% at 0.75 ppm **Bronchospasm:** In sensitive individuals	**Systematic Review:** 67 studies showing no threshold **Asthmatic Sensitivity:** Bronchoconstriction at 0.25-0.75 ppm **Mechanistic Review:** Oxidative stress pathways
Heart Failure: Increased risk **Arrhythmias:** 2× higher rates **Neurological Sequelae:** 15-50% of survivors **Cognitive Decline:** At 3 ppm increase **Depression:** >40% incidence post-exposure **Tissue Hypoxia:** COHb formation	**Framingham Heart Study:** Exhaled CO and heart failure **Taiwan Cohort:** 20,942 exposed showing cardiac effects **Brain Imaging:** Reduced blood flow at low levels
Leukemia (AML): 2-3× risk **Blood Disorders:** At 1 ppm exposure **DNA Damage:** Chromosomal aberrations **Sperm Abnormalities:** Aneuploidy **No Safe Threshold:** Group 1 carcinogen	**Pliofilm Workers Study:** Established leukemia link **Chinese Benzene Cohort:** 74,828 workers showing dose-response **Meta-Analysis:** Quantified AML risk **Low-Level Effects:** Blood effects at 1 ppm
IQ Loss: 7.4 points <10 μg/dL blood **CV Mortality:** 70% increase **412,000 US deaths/year** attributable to historical exposure to leaded gas **Kidney Disease:** Progressive **Hypertension:** Dose-dependent **No Safe Threshold**	**IQ Meta-Analysis:** 1,333 children, no threshold **NHANES Mortality:** 14,289 adults, CV deaths **Rochester Study:** Effects <10 μg/dL **Population Impact:** 170M Americans exposed

Biological and Particulate Contaminants

POLLUTANT	TYPICAL INDOOR LEVELS	MAJOR SOURCES
Gram-Negative Bacteria (Endotoxins)	**Typical:** 10-1000 CFU/m³ **Endotoxin in PM2.5:** 0.13-0.17 EU/m³ **Netherlands Limit:** 90 EU/m³ (8-hr) **European Guideline:** <50 EU/m³ *Health effects at 28 EU/m³*	• Water-damaged materials • Contaminated HVAC systems • Pet presence • Flooding/moisture • Food waste
Gram-Positive Bacteria	**Winter Homes:** 19-607 CFU/m³ **Schools:** 924-2750 CFU/m³ **Mean:** 212 CFU/m³ *Varies with occupancy*	• Human skin/respiratory shedding • Textiles and soft furnishings • Dust reservoirs • Air handling systems
Mycotoxin-Producing Molds: Stachybotrys (Black Mold)	**When Detected:** <32 counts/m³ **Detection Threshold:** Any level warrants investigation **Water Activity:** >95% required *Indicates severe water damage*	• Water-damaged gypsum board • Chronic leaks • Flooding • High humidity (>95%) • Cellulose materials
Mycotoxin-Producing Molds: Aspergillus	**Indoor Median:** 82 CFU/m³ **Outdoor Median:** 540 CFU/m³ **Combined Asp/Pen:** <700 counts/m³ **AIHA Guideline:** <250 CFU/m³ total fungi	• Damp building materials • House dust • Stored grains • Compost • HVAC systems
Mycotoxin-Producing Molds: Penicillium	**Typical with Aspergillus:** <700 counts/m³ **Indoor Median:** 82 CFU/m³ **Elevated in damp homes**	• Water-damaged materials • Food spoilage • Indoor plants • Carpets • Wallpaper
Non-Mycotoxin Molds: Cladosporium	**Summer Outdoor:** 2,000-50,000 spores/m³ **Winter:** Very low **Symptom Threshold:** 3,000 spores/m³ **Canadian Guideline:** <500 CFU/m³ *Most widespread allergenic mold*	• Outdoor infiltration • Damp gypsum board • Painted walls • Wood surfaces • Carpets • HVAC components
Non-Mycotoxin Molds: Alternaria	**Symptom Threshold:** 100 spores/m³ **Peak Season:** Warm, dry weather **Indoor:** Variable with outdoor *10× more potent than Cladosporium*	• Outdoor spores entering homes • Damp, poorly ventilated spaces • Carpets • Bedding • Textiles
Gypsum/Drywall Dust	**OSHA Limit:** 50 μg/m³ (8-hr respirable) **Sanding:** Up to 150 mg/m³ **Particle Size:** ~10 microns *Contains calcium sulfate, talc, mica*	• Drywall installation • Sanding operations • Demolition • Deteriorating wallboard • Renovations
Pet Dander	**Fel d 1 (cat):** Detectable in all homes **Can f proteins (dog):** Widespread **Airborne Duration:** Hours **HEPA Reduction:** 30-40% *15% Americans allergic*	• Cats (Fel d 1 protein) • Dogs (Can f proteins) • Transfer on clothing • Settled in furniture • Carpets and bedding

HEALTH EFFECTS	KEY SCIENTIFIC EVIDENCE
Illness Episodes: +0.32/year in children **Acute Airway Obstruction** **Hypersensitivity Pneumonitis** **Chronic Bronchitis** **Fever, Cough, Dyspnea** **Inflammatory Response:** TNF-α, IL-1β	**Children's Health Study:** Illness correlation **Indoor Endotoxin Study:** Urban concentration patterns **Netherlands Health Council**: Occupational exposure limits
Respiratory Infections **Allergic Responses** **Skin Conditions** (Staph aureus) **Indoor Air Quality Complaints** **Opportunistic Infections**	**Indoor Microbiome Studies:** Human contribution **Seasonal Variation Studies:** Winter elevation patterns **Occupancy Studies:** Direct correlation with human presence
Cognitive Impairment **Chronic Fatigue** **Nosebleeds** **Memory Problems** **Produces:** Satratoxins G & H	**Cleveland Outbreak:** Infant pulmonary hemosiderosis **Mycotoxin Detection:** Airborne confirmation **Indoor Air Study:** Health associations
Lung Infections (immunocompromised) **Allergic Aspergillosis** **Asthma Exacerbation** **Produces:** Aflatoxins, ochratoxins **Cancer Risk** (aflatoxin B1)	**A. fumigatus Studies:** Invasive aspergillosis in hospitals **Mycotoxin Analysis:** Indoor detection methods **WHO Guidelines:** Health-based recommendations
Childhood Asthma Development **Allergic Rhinitis** **Produces:** Ochratoxin A, patulin **Kidney Toxicity** (ochratoxin) **Immune Suppression**	**Boston Children Study:** Asthma link **Cincinnati Study:** Wheeze association **European Studies:** Ochratoxin A in indoor dust
Allergic Rhinitis **Asthma Attacks** **Nasal Congestion** **Eye Irritation** **Sneezing** **Major Allergens:** Cla h 1, Cla h 2	**Global Allergen Studies:** Most common outdoor mold **Threshold Studies:** 3,000 spores/m^3 for symptoms **Canadian Housing Studies:** Prevalence and guidelines
Main Fungal Asthma Cause **Severe Asthma** association **Hay Fever** **Alt a 1 Sensitization:** 9.5-20% population **Poor Asthma Prognosis**	**European Allergy Study:** 9.5% sensitization rate **Beijing Study:** Most common aeroallergen **Severity Studies:** Associated with severe asthma
Respiratory Irritation **Eye/Nose/Throat Irritation** **Coughing** **Allergic Reactions** **Silicosis Risk** (if silica present) **Lung Function Decline**	**NIOSH Studies:** Exposure during sanding **Occupational Health Reviews:** Long-term lung effects **Construction Worker Studies:** Chronic respiratory symptoms
Asthma Episodes: 3 in 10 asthmatics **Allergic Rhinitis** **Eye Irritation/Watering** **Skin Rashes** **Wheezing/Coughing** **20% adults, 30% children affected**	**Allergen Distribution Studies:** Present even in pet-free homes **HEPA Filtration Studies:** 30-40% reduction efficacy **Epidemiological Studies:** 15% population allergic

INDOOR AIR QUALITY:

Biological and Particulate Contaminants

POLLUTANT	TYPICAL INDOOR LEVELS	MAJOR SOURCES
Pollen	**Variable:** 0-2000+ grains/m³ **Peak Seasons:** Spring (tree), Summer (grass), Fall (ragweed) **Indoor:** 10-70% of outdoor *25 million Americans affected*	• Window/door infiltration • Ventilation systems • Transfer on clothing/hair • Pet fur • Open windows
Dust Mites	**Sensitization Risk:** ≥2 μg/g dust **Asthma Risk:** ≥10 μg/g dust **Typical Homes:** 0.2-14 μg/g **Thrives:** >50% humidity *Der p 1 and Der f 1 allergens*	• Mattresses/pillows • Bedding • Carpets • Upholstered furniture • Human skin scales (food)

The Inflammation Cascade Explained

Think of inflammation as your body's defense system. Acute inflammation (like when you cut yourself) is good - it heals you. But chronic inflammation from breathing bad air is like having your immune system constantly activated at 30-40% intensity, slowly damaging everything.

C-REACTIVE PROTEIN (CRP) - YOUR BODY'S GENERAL FIRE ALARM

- What it is: A protein your liver makes when there's inflammation anywhere in your body
- Normal range: <1.0 mg/L is optimal, 1-3 is average, >3 is high risk
- 40% elevation means: If your baseline is 1.0, bad air pushes you to 1.4 - moving you toward the danger zone
- Why biohackers care: High CRP correlates with heart disease, cancer, depression, and accelerated aging
- You might know this from: Your annual blood work or cardiovascular risk panels

IL-6 (INTERLEUKIN-6) - THE AGING ACCELERATOR

- What it is: A signaling molecule that tells your body to create more inflammation
- The biohacker connection: This is what spikes after poor sleep, intense exercise, or stress
- 25% increase means: Your body ages faster, recovers slower, and stays inflamed longer
- Downstream effects: IL-6 triggers CRP production, disrupts sleep (it's highest at night), and impairs muscle recovery
- Why it matters: Centenarians have unusually low IL-6 levels - it's a longevity biomarker

HEALTH EFFECTS	KEY SCIENTIFIC EVIDENCE
Per 10 grains/m³ increase: • Lower respiratory: +1% • Any allergic symptom: +2% • Upper respiratory: +7% • Ocular symptoms: +11% **Annual Cost:** >$3 billion	**Systematic Review:** Dose-response relationships **Economic Studies:** $3 billion annual costs **Indoor Infiltration Studies:** Building characteristics impact
3× Asthma Risk at age 7 **5× Late-onset Wheeze** **Allergic Rhinitis** **Eczema Exacerbation** **Bronchial Hyperresponsiveness** **OR 1.94 for asthmatics**	**Threshold Studies:** 2 μg/g sensitization **Birth Cohort Studies:** 10 μg/g asthma threshold **Meta-Analyses:** Consistent asthma associations





TNF-α (TUMOR NECROSIS FACTOR ALPHA) – THE AUTOIMMUNE TRIGGER

- What it is: A master inflammatory cytokine that can actually kill cells (hence "necrosis")
- 30% elevation means: Your immune system becomes more likely to attack your own tissues
- The cascade effect: TNF-α triggers other inflammatory molecules, creating a vicious cycle
- Real world impact: This is what drugs like Humira and Enbrel block in autoimmune conditions
- For biohackers: Elevated TNF-α = brain fog, joint pain, fatigue, and increased risk of developing autoimmune conditions

OXIDATIVE STRESS – YOUR CELLULAR RUSTING

- What it is: Free radicals (unstable molecules) damaging your cells faster than your antioxidants can repair them
- 45% higher means: Your cells are aging 45% faster than they should
- The analogy: Like leaving metal in rain – it rusts. Your cells are "rusting" from the inside
- Measurable via: 8-OHdG in urine, F2-isoprostanes, or glutathione/glutathione disulfide ratio
- Biohacker relevance: This is why people take NAD+, glutathione, and antioxidants

How Air Quality Triggers This Cascade

When you breathe mold spores, PM2.5, or VOCs, here's the sequence:



1. Immediate: Lung cells detect foreign particles and release IL-6
2. Hours: IL-6 signals liver to produce CRP
3. Days: TNF-α levels rise as immune system stays activated
4. Ongoing: Oxidative stress accumulates as your antioxidant systems get overwhelmed
5. Result: Chronic low-grade inflammation that underlies virtually every disease

Why These Specific Percentages Matter

In biohacker terms, these increases are like:

- Running your HRV 20 points lower constantly
- Having your recovery scores permanently in the yellow/red
- Aging at 1.3x speed instead of 1x
- Giving yourself the inflammation equivalent of eating inflammatory foods at every mealeam

EVERY BREATH MATTERS

4.14 Million

Estimated global deaths annually from PM 2.5 alone

824 Million

IQ points lost in the US cumulatively from childhood lead exposure

4.2 Years

Lifespan extension opportunity from average US outdoor levels to pristine air quality

COMPLIED FROM

2,000

peer-reviewed studies

50+

landmark cohorts

30+

meta-analyses

How This Guide Was Built
This reference synthesizes evidence from:

- **New England Journal of Medicine (NEJM)**
- **Journal of the American Medical Association (JAMA)**
- **The Lancet & The Lancet Public Health**
- **Environmental Health Perspectives**
- **Nature & Science**
- **American Journal of Respiratory and Critical Care Medicine**
- **EPA Integrated Science Assessments**
- **WHO Air Quality Guidelines**

Methodology: Every health effect quantified represents findings replicated across multiple independent research teams, in different populations, using varied methodologies, yet reaching consistent conclusions.

Last Updated: November 2025
Version: 2.1

Scientific Advisory
Lution Labs Clinical Research Team

+ lution

Contact
science@lutionlabs.com

This guide provides scientific education, not medical advice. Consult healthcare providers for personal health decisions.

Measuring What You Breathe.
Optimizing Everything.